

General background text Pharmacogenetics – Human Leukocyte Antigen (HLA)

Last updated: 1 March 2017

Definitions in pharmacogenetics

The **genotype** is the hereditary information about a specific characteristic of an individual. This information is located in the genes, in the DNA that consists of nucleotides. The piece of the DNA that carries information for one specific hereditary characteristic is called a **gene**. The DNA is divided into chromosomes, which usually occur in pairs. A person generally has two copies (**alleles**) of a gene, one on each of the chromosomes of a chromosome pair.

The **phenotype** indicates what the final manifestation (phenotypic state) of a certain genotype is. This can involve the functionality of a protein (for example the enzyme or the receptor), but also the physical manifestation of a disease. The phenotype is a result of the genotype that a person possesses, the degree of expression of the gene in question and the combination with environmental factors such as co-medication, diet and disease conditions.

Variations can exist in a population within the DNA that encodes a protein. Variations can result in alleles that encode for proteins with no or reduced activity. The simplest form of variations are “**single-nucleotide polymorphisms**” (**SNPs**), in which a certain part of a gene differs by only one nucleotide. If a gene variation occurs in at least 1% of the population, then this is referred to as a genetic **polymorphism**. There can be a number of different polymorphisms for a certain allele.

HLA proteins

Human Leukocyte Antigen (HLA) is the name for proteins of the human Major Histocompatibility Complex (MHC). These proteins are divided into 2 classes: class I and class II, each with their own function.

MHC class I proteins are on the surface of almost all body cells. They bind peptides in the cell and present them on the surface of the cells allowing immune cells to ‘see’ whether the cell produces proteins that do not belong in the cell (e.g. after a viral infection). Any cells presenting abnormal peptides are killed.

MHC class II proteins are only expressed on the surface of specialised immune cells. They bind peptides from proteins that have been internalised by cells from the outside. They present these peptides on the surface allowing immune cells to ‘see’ whether any non-self proteins are present in the body. Any cells presenting non-self peptides cause activation of the immune system.

Class I and class II proteins are each encoded by 3 different polymorphic genes (HLA-A, HLA-B, HLA-C and HLA-DP, HLA-DQ and HLA-DR respectively). The number of different alleles per polymorphic gene is very high, so that the chances of two people with the same combination of HLA alleles meeting is very low (with the exception of identical twins).

HLA polymorphisms and clinical consequences

The polymorphic HLA proteins differ in peptides they can bind and therefore determine the nature of the peptides the body can mount an immune response to. Pathogens consist of multiple proteins, each of which are broken down to a multitude of peptides. People with different polymorphic variations in HLA can therefore all mount immune responses to pathogens and other non-self proteins, but the intensity of the response may differ between individuals.

HLA polymorphisms may therefore play an important role in the development of immune responses or antibodies to protein medicines or to medicines that lead to modification of proteins in the body (e.g. by medicines or metabolites binding to proteins).

As the immune system responds more strongly to pathogens/proteins it has encountered before, it is an important property of immune-mediated hypersensitivity that the hypersensitivity reaction becomes more severe with longer usage or rechallenge with the medicine.

Genotyping

The process of genotyping is used to determine the genotype. It indicates which alleles of the various HLA genes are present in the tested individual. Each allele has a name that consists of the name of the gene, followed by a star (*) and a number, an example of a possible HLA allele is HLA-B*5701.

As the activity of HLA alleles is highly specific, only one allele of the 6 polymorphic genes generally has a distinct effect on protein-based and other medicines. In the event of HLA drug interactions, only one set of alleles of a polymorphic gene therefore generally needs to be determined.

As HLA alleles are determined as part of tissue typing to establish whether donors and recipients are transplant compatible, typing of polymorphic HLA alleles is a standard procedure in many laboratories of immunology and clinical chemistry. However, this generally uses phenotyping rather than the more expensive genotyping. The standard genotyping method is DNA sequence-based typing.

There are cheaper methods available for specific alleles, such as PCR with sequence-specific primers for the identification of HLA-B*5701. HLA-DRB typing is also possible using an HLA chip.

Phenotyping

The process of phenotyping is used to determine the phenotype, which means: establishing which HLA proteins are present. This can be done using standard serology tests, involving staining cells using antibodies against various HLA proteins. Another option is functional testing (complement-dependent cytotoxicity tests for identification of class I proteins). A disadvantage of these tests is that they cannot distinguish between subtypes of polymorphic HLA proteins, meaning that they can be used to determine whether a person has HLA-B57, but not whether this person has HLA-B*5701, HLA-B*5702 or any of the other subtypes.

Ethnic variation in prevalence of phenotypes and allele frequency

The frequency of occurrence of the various HLA alleles varies significantly between ethnic groups.

The allele frequencies of six pharmacogenetic implicated HLA alleles identified in various populations are given in table 1.

Table 1. Ethnic variation in allele frequency (% of people with this allele)

Ethnicity	Country	HLA-B*5701	HLA-B*44	HLA-A*3101	HLA-B*1502	HLA-B*1511	HLA-B*5801
Caucasian		4.2 - 7.6	6.7 - 14.9	3.0-13.4	0-1.2	0-0.38	0-4.5
	The Netherlands	3.4	12.4	3.0-6.6	0	0	0.6
	Northern Europe			3.0-6.6	0-0.04	0-0.02	0-0.9
	Southern Europe	1-4		3.6-5.0	0-1.2	0	0-4.5
	Mediterranean	1-2					
Asian	USA	1.0	5.2				
	China	0		0.8 -12.6	0-59	0-24	2.3-17
	Taiwan			5.6	8.4-12	1.8	9.8-10.1
	Japan	0		16.8	0.06-0.2	0.8-1.8	0.4-0.58
	Thailand	4-10		2.6	17	2.0	6.1-8.4
	Malaysia			0.8-2.6	0.04-31	0.22-0.52	5.8-14.0
	India	5-20			0-12		5.8-8.8
	South Korea			11.2	0.4-4.4	3.3-4.0	6.08-6.5
	Vietnam				26		6.5
	Myanmar			4.6	18-34		0.91-4.17
	Indonesia				21-31		4.2-6.0
	Philippines				39	0	
	Middle East	1-2			0-1.2	0	1.4-6.8
African	Sub Saharan	0.2 - 3		0.8-1.8	0	0	0.7-15
	Northern Africa			3.0	0	0	1.4-4.0
African-American		2.4 - 9.0	5.5	2.0	0-0.4	0	
Hispanics		1.9 - 4.7	10.9				1.1-1.45
Indians		2.1	7.8				
South				5.6-44	0-1.6	0-1.0	0.50-2.77

American							
----------	--	--	--	--	--	--	--

Literature

1. Martin AM et al. HLA-B*5701 typing by sequence-specific amplification: validation and comparison with sequence-based typing. *Tissue Antigens* 2005;65:571-4.
2. Jiao J et al. Hepatitis C virus genotypes, HLA-DRB alleles and their response to interferon-alpha and ribavirin in patients with chronic hepatitis C. *Hepatobiliary Pancreat Dis Int* 2005;4:80-3.
3. Cao K et al. Analysis of the frequencies of HLA-A, B, and C alleles and haplotypes in the five major ethnic groups of the United States reveals high levels of diversity in these loci and contrasting distribution patterns in these populations. *Hum Immunol* 2001;62:1009-30.
4. Waters LJ et al. Prospective HLA-B*5701 screening and abacavir hypersensitivity: a single centre experience. *AIDS* 2007;21:2533-4.
5. Nolan D et al. Pharmacogenetics: a practical role in predicting antiretroviral drug toxicity? *J HIV Ther* 2003;8:36-41.
6. Phillips EJ. Genetic screening to prevent abacavir hypersensitivity reaction: are we there yet? *Clin Infect Dis* 2006;43:103-5.
7. Abel S et al. Abacavir hypersensitivity. *N Engl J Med* 2008;358:2515.
8. Sadiq ST et al. Uncertainties of routine HLA B*5701 testing in black African HIV cohorts in the UK. *Sex Transm Infect* 2007;83:181-2.
9. Schipper RF et al. HLA gene haplotype frequencies in bone marrow donors worldwide registries. *Hum Immunol* 1997;52:54-71.
10. Allele Frequency Net Database: <http://www.allelefreqencies.net>